



# International Journal of Homoeopathic Sciences

E-ISSN: 2616-4493

P-ISSN: 2616-4485

[www.homoeopathicjournal.com](http://www.homoeopathicjournal.com)

IJHS 2023; 7(4): 414-422

Received: 25-10-2023

Accepted: 27-11-2023

**Ankitha N Karanik**

Principal Investigator, Final  
year BHMS, Father Muller  
Homoeopathic Medical  
College, Mangaluru,  
Karnataka, India

**Dr. Sneha Jnanakshee HR**

PG Scholar, Department of  
Organon of Medicine and  
Homeopathic Philosophy,  
Father Muller Homoeopathic  
Medical College, Mangaluru,  
Karnataka, India

**Dr. Skandhan S Kumar**

BHMS MD Organon of  
Medicine (Gold medallist),  
Assistant Professor,  
Department of Organon of  
Medicine and Homoeopathic  
Philosophy, Father Muller  
Homoeopathic Medical  
College, Mangaluru,  
Karnataka, India

**Corresponding Author:**

**Ankitha N Karanik**

Principal Investigator, Final  
year BHMS, Father Muller  
Homoeopathic Medical  
College, Mangaluru,  
Karnataka, India

## To determine the potential of agrohomoepathy by using calcarea phosphorica 200c by seed priming technique and analyzing the germination rate and growth rate of *Vigna unguiculata* plant (cow pea) plant

**Ankitha N Karanik, Dr. Sneha Jnanakshee HR and Dr. Skandhan S  
Kumar**

**DOI:** <https://doi.org/10.33545/26164485.2023.v7.i4f.1007>

### Abstract

This study was undertaken to know the effectiveness of calcarea phos 200C on Agrohomoepathy, a single blinded study that shows experimental group had good yield when compared to control group.

**Background:** This article gives an insight to common man who is interested in farming or agriculture, to get accustomed with new technique and effectiveness of homoeopathic medicines in Agrohomoepathy, how we can increase germination rate and growth rate of various plants cost effectively, so that we can take various common problems that arises during cultivation of such plants.

**Methods:** Various old literatures throws light upon Agrohomoepathy, and its role on homoeopathic medicine upon different plants. New techniques like Seed Priming is highlighted in this present studies.

**Conclusion:** Gives idea to a common man regarding effectiveness of homoeopathic medicine in increasing the quality of yield by improving germination and growth rate without using synthetic chemicals.

**Keywords:** *Vigna unguiculata* plant, calcarea phosphoricum, centesimal potency, seed priming technique, germination and growth rate, agro homoeopathy

### Introduction

Agriculture is the backbone of the Indian economy; more than 50% of Indian people's livelihood directly depends on it, but contributes only 18% of nations GDP. So, India plays great emphasis on natural approach and is willing to expand resources on agro homeopathy and produce applicable results for this country.

Agro homeopathy is a branch of science which deals with using homoeopathic medicine on plants influencing its growth, metabolites production, essential yield, photochemical process and nutritive values when applied on germinating seeds or on growing plants.

The idea of testing the Homoeopathic remedies on plants was invented by Dr. Baron von Boenninghausen. He has quoted in his writings that the medicines which were left out were thrown to his plant pots and observed effect on plants.

Homoeopathic preparation can be applied on the soil, leaves or any part of the plant to increase, decrease the production of bioactive substances and directly increase its quality.

Modern agriculture practices with an intensive approach (e.g., heavy tillage, synthetic agrochemicals, hybrid varieties and mono cropping) may further deteriorates fragile lands by accelerating the loss of organic matter, degradation of soil and environmental damage (deterioration of water and air quality, loss of biodiversity, and biomagnifications) thus, several attempts have been carried out to achieve ecosystem stability. Vital aspects of crop improvement through maintaining sustainability can be accomplished by modulating the metabolism of seed that would be achieved by seed priming technique.

### What is seed priming technique?

Seed priming is pre sowing seed treatment that allows controlled hydration of seeds to imbibe water and go through the first stage of germination but does not allow radical protrusion through the seed coat.

The aim is to develop natural, simple and inexpensive methods to increase agricultural production without using synthetic chemicals, so that a wide range of agricultural hazards can be controlled. This study is an affordable therapeutic treatment used in cultivation of plants. With an increase in the number of farmers and researchers are opting for natural and organic methods of farming this study would be useful. With the development in the field of agriculture, there is increased in organisms affecting the plants, hence using of homoeopathic in agriculture as a conventional agriculture way to maintain agro-ecological balance in this field is essential.

Article by department of food science and technology, university of reading, UK States that optimization of germination process of cowpea by response surface methodology is 8 hours with germination time as 52 hrs, hence this research is done as a part of continued study.

### Why Calcareo phosphorica?

Calcareo phosphorica is one of a twelve tissue salts Proved by - Schussler

Scientific name is Phosphate of lime

Common name is Calcium Phosphate

Calcium and phosphate is a vital component of soil required for the plant growth

Soil studies show that calcium and phosphate was less in Mangalore soil hence we have chosen this medicine

### Why in centesimal potency?

Dr Borieck Material Medica says Dose – First to third trituration. Higher potencies often more effective.

Agro homeopathy repertory says

**Leaves; MARGINS;** yellow- calcarea phos

**Leaves; Dry-** calcarea phos

**Leaves; Leathery-** calcarea phos

**Leaves; Rolled-** calcarea phos

**Leaves; Wrinkled-** calcarea phos

**Named Diseases; Mildew-** calcarea phos

**Named Diseases; Sooty-** calcarea phos

**General; Brittle-** calcarea phos

**General; Discoloration;** brown- calcarea phos

**General; Straggly;** calcarea phos

**Generative; Pollination;** excessive- calcarea phos

**Generative; Stamen;** long- calcarea phos

**Fruits; Rotting-** calcarea phos

**Epidermis; Cracks-** calcarea phos

**Epidermis; Thin-** calcarea phos

**Study Design:** 40 seeds of cow pea will be selected after certified by Botanist.

20 seed will be soaked in 2 jars respectively for a period of 8 hrs for seed priming technique.

Coder will be assessing the jars, medicine and hence principal investigator or guide is unaware of the procedure.

20 seeds will be put in one jar and medicine Calcareo Phosphorica 200 C (1 drop of medicine with 50 ml of distilled water) will be introduced in one jar and labeled as S1 (Experiment group).

20 seeds will be put in one jar and medicine (1 drop of rectifying spirit with 50ml of distilled water) will be introduced in another jar and labeled as S2 (control group).

After 8 hrs the seeds will be shifted to respective grow bags filled with soil and drainage holes. Regular monitoring will be made for a period of 10 days (to avoid insects or to regulate water content and to regulate temperature) and will be labeled as S1, S2 respectively.

Regular watering will be done morning and evening, everyday with normal water and factors like germination and growth rate will be recorded for 10 days.

On 11 th day the germinated seed shoot length and root length will be measured using thread and scale and will be assessed for germination rate. And these seeds will be introduced into their respective grow bags and will be named as S1a, S1b, S1c, and S1d.... S2a, S2b, S2c, S2d... regular watering and monitoring will be done for at respective time intervals for a period of 3 months.

At the end of 3 months plants will be removed from the soil, weight and shoot length will be measured and considered to be fresh weight.

For dry weight the plants are then kept in hot air oven to and dried at 110°F for 5hrs at and weight will be measured.




### The observations are as followed

**Table1:** S1 Experiment group data's in centimeters on respective dates. (SHOOT in CMS)

S1(days)	1	2	3	4	5	6	7	8	9	10	11	1 Month	2 Month	3 Month
A		1	3	14	15	16	18	18.2	19	21	23	60	98	149
B		2	4	12	16	17	18	18.3	19	22	23	117	202	308
C		1.5	3	10	12	15	16	17.8	18	20	22	80	139	249
D		3	6	15	16	17	18	19	22	23.5	24.7	115	230	410
E		2	5	14	15	16	18	20	21	22	23	110	200	400
F		1	3	8	9	10	11	11.7	15	19	22	86	112	203
G		1	3	7	10	10.8	11	11.9	15.3	18	20	87	115	205
H		2.1	3	7.5	10	11	12	15	18	21	22	122.5	220	387
I		1.7	2	8.5	11	12	13.5	14	16	18	21	60	78	100
J		1	2	6	11	13	15	15.8	16	18	20	78	130	212
K		2	3	6	12	12.7	15	18	19	20	21.5	40	60	88
L		1	2	5	11	12	13	15	18	20	22	40	68	84
M		2	4	12	13	14	15	18	19	21	22.7	88	122	217
N		0	3	13	14	15	16	19	19.7	22	23	90	150	273
O		0	4	11	13	15	16	19	20	21	23	60	100	168
P		0	4	10	13	16	18	19	20	22	24	125	206.5	306
Q		0	4	9	12	14	17	19	20	21	23	97.5	198	259
R		0	3	6	9	10	10.2	12	13	18	19	48	57	79
S		0	0	7	9	9.5	10	10.5	14	20	21	47.5	55	70
T		0	0	0	3	6	7	12.5	19	20	21	45	50	66

**Table 2:** S2 Control group data's in centimeters on respective dates. (SHOOT in CMS)

S2	1	2	3	4	5	6	7	8	9	10	11	1 Month	2 Month	3 Month
A	1	2	6	7	8	11	13	15	20	21		82	126	200
B	0.5	2	6	8	10	13	15	18	19	20		86	133	176
C	2	4	13	14.5	15	17	18	21	21.5	22.5		110	191	245
D	1	1.5	5	8	12	15	16	16.5	18	20		49	91	117
E	1	2	10	12	13	14	14.5	16	17	20		52	100	129
F	1.7	2	6	9	11	12	12.7	13	16	20		30	56	100
G	0	3	5	9	11.5	12	12.2	13	15	19		25	35	45
H	0	4	8	12	12.7	14	15	18	21	22		86	123	210
I	4	4	8	12	13	15	17	19	21	22		36.2	53	70
J	0	0	11	11.5	15	16	16.5	17.8	20	22.7		42	98	113
K	0	0	10	11.5	14	15	16	17.7	20	22.5		30	51	68
L	0	0	7	12	14	14	15.5	16	17	19		29	59	73
M	0	0	7	11	13	13	15	16.2	17	19		30	60	78
N	0	0	6	11	12.6	13	15	17	18	20		26	38	49
O	0	0	6	10	12.8	15	16	18	20	21		28.5	35	0
P	0	0	6	10	13	14	16	18	20	22		28	32	0
Q	0	0	5	9	10	11	14	17	19	21		24	0	0
R	0	0	0	0	12	12.5	13	15	18	21		23	0	0
S	0	0	0	0	0	0	0	0	16	19		0	0	0
T	0	0	0	0	0	0	0	0	17	20		0	0	0








**Fig 1:** Shows the seed soaking, germination and growth of the plant

**Table 3:** Length of the root in centimeters on 11<sup>th</sup> day and 3 months (ROOT in CMS)

S1	11 <sup>th</sup> Day	3 Month	S2	11 <sup>th</sup> Day	3 Month
A	30	12.5	A	20	11
B	32	10	B	29	22
C	44	5.5	C	30	16
D	28	15	D	24	5
E	24	11	E	10	15
F	30	5	F	15	5
G	27	12	G	15	7
H	30	6	H	15	13
I	26	11	I	15	10
J	23	4	J	28	13
K	21	23	K	21	7
L	30	21	L	25	7
M	23	7.5	M	15	5
N	17	8	N	16	7
O	26	8	O	17	
P	32	9	P	9	
Q	34	8	Q	14	
R	20	10	R	22	
S	32	11	S	21	
T	26	8	T	19	




**Fig 2:** Shows the root length of the plant

**Table 4:** Dry weight and wet weight of the plant Number of pods by the end of 3rd month

	S1	S2
Fresh weight	241.4	109.7
Dry weight	54.4	18.9
No of pods		
S1	14	
S2	1	





**Fig 3:** Shows the wet weight and dry weight of both the group. Also shows the pod.

**Statistical analysis and results****Table 5:** Statistical data

<b>Tests of Normality</b>					
	Group	Kolmogorov-Smirnov <sup>a</sup>		Shapiro-Wilk	
		Statistic	Sig.	Statistic	Sig.
q11	Experiment	.150	.200*	.956	.467
	Control	.263	.010	.861	.031
Month1	Experiment	.156	.200*	.927	.136
	Control	.200	.134	.825	.010
Month2	Experiment	.170	.134	.910	.065
	Control	.225	.054	.904	.128
Month3	Experiment	.140	.200*	.928	.140
	Control	.169	.200*	.903	.125

**Table 6:** Statistical data

Descriptive				
	Group		Statistic	
q11	Experiment	Mean	22.0450	
		Median	22.0000	
		Std. Deviation	1.41402	
		Minimum	19.00	
		Maximum	24.70	
		Range	5.70	
		Interquartile Range	2.00	
	Control	Mean	20.6929	
		Median	20.0000	
		Std. Deviation	1.38867	
		Minimum	19.00	
		Maximum	22.70	
		Range	3.70	
	Interquartile Range		2.38	
Month1		Experiment	Mean	79.8250
			Median	83.0000
			Std. Deviation	28.64404
			Minimum	40.00
			Maximum	125.00
			Range	85.00
			Interquartile Range	55.88
		Control	Mean	50.9429
Median			39.1000	
Std. Deviation			28.15478	
Minimum			25.00	
Maximum			110.00	
Range			85.00	
Interquartile Range			53.25	
Month2		Experiment	Mean	129.5250
			Median	118.5000
			Std. Deviation	60.90361
			Minimum	50.00
			Maximum	230.00
			Range	180.00
			Interquartile Range	129.00
		Control	Mean	86.7143
Median			75.5000	
Std. Deviation			44.79428	
Minimum			35.00	
Maximum			191.00	
Range			156.00	
Interquartile Range			71.25	
Month3		Experiment	Mean	211.6500
			Median	208.5000
			Std. Deviation	112.40307
			Minimum	66.00
			Maximum	410.00
			Range	344.00
			Interquartile Range	206.75
		Control	Mean	119.5000
Median			106.5000	
Std. Deviation			64.23485	
Minimum			45.00	
Maximum			245.00	
Range			200.00	
Interquartile Range			112.50	

**Table 7:** Statistical data

<b>Group Statistics</b>					
	Group	N	Mean	Std. Deviation	t
q11	Experiment	20	22.045	1.414	3.206
	Control	20	20.685	1.265	p=0.003 hs
Month1	Experiment	20	79.825	28.644	3.811
	Control	18	45.372	26.880	p<0.001 vhs
Month2	Experiment	20	129.525	60.904	2.699
	Control	16	80.063	45.494	p=0.011 sig
Month3	Experiment	20	211.650	112.403	2.760
	Control	14	119.500	64.235	p=0.009 hs

**Table 8:** Statistical data

Descriptive						
q11						
Group		N	Mean	Std. Deviation	Minimum	Maximum
Experiment	Day 11	20	22.045	1.414	19.000	24.700
	One month	20	79.825	28.644	40.000	125.000
	Two months	20	127.475	58.115	50.000	230.000
	Three months	20	211.650	112.403	66.000	410.000
	Total	80	110.249	94.503	19.000	410.000
Control	Day 11	20	20.685	1.265	19.000	22.700
	One month	20	55.835	44.364	23.000	202.000
	Two months	14	73.000	44.232	32.000	191.000
	Three months	14	119.500	64.235	45.000	245.000
	Total	68	62.138	54.551	19.000	245.000

**Table 9:** Statistical data

<b>ANOVA</b>			
q11			
Group		F	p
Experiment	Between Groups	30.548	<0.001 vhs
Control	Between Groups	15.177	<0.001 vhs

**Post Hoc Tests****Table 10:** Statistical data

<b>Multiple Comparisons</b>				
Dependent Variable: q11				
Bonferroni				
Group	(I) period	(J) period	Mean Difference (I-J)	p
Experiment	Day 11	One month	-57.780	.037 sig
		Two months	-105.430	<0.001 vhs
		Three months	-189.605	<0.001 vhs
	One month	Two months	-47.650	.137
		Three months	-131.825	<0.001 vhs
		Two months	-84.175	<0.001 vhs
Control	Day 11	One month	-35.150	.068
		Two months	-52.315	.005hs
		Three months	-98.815	<0.001 vhs
	One month	Two months	-17.165	1.000
		Three months	-63.665	<0.001 vhs
		Two months	-46.500	.032 sig

<b>Group Statistics</b>				
	Group	N	Mean	Std. Deviation
Length of the root- - 11thday	Experiment	20	27.750	5.928
	Control	20	19.000	5.982
Length of the root- - 3 months	Experiment	20	10.275	4.852
	Control	14	10.214	5.071

**Table 11:** Statistical data

<b>Independent Samples Test</b>		
	t-test for Equality of Means	
	t	p
Length of the root- - 11thday	4.646	<0.001 vhs
Length of the root- - 3 months	.035	.972






**Table 12:** Statistical data

Paired Samples Statistics				
Group		N	Mean	Std. Deviation
Experiment	Length of the root- - 11thday	20	27.750	5.928
	Length of the root- - 3 months	20	10.275	4.852
Control	Length of the root- - 11thday	14	19.857	6.371
	Length of the root- - 3 months	14	10.214	5.071


**Table 13:** Statistical data

Paired Samples Test						
Group			Paired Differences		t	p
			Mean	Std. Deviation		
Experiment		Length of the root- - 11thday - Length of the root- - 3 months	17.475	8.320	9.393	<0.001 vhs
Control		Length of the root- - 11thday - Length of the root- - 3 months	9.643	6.380	5.655	<0.001 vhs

**Fig 4:** Graphical representation of length of the root of both experiment group and control group**Fig 5:** Graphical representation of lenth of the root of both experiment group and control group



**Fig 6:** Shows the graphical representation of fresh weight of both experiment and control group



**Fig 7:** Shows the graphical representation of no of pods by the end of third month of both experiment and control group

## Discussion

The study which is conducted using plant has more effective study. Where the P value is greater than .05

## Results: p value Effective study

The P value is  $p < 0.5$  experiment group is proved to be effective than control group.

## Conclusion

This study suggest that Agrohomoepathy is less harmful than conventional method of farming which is practiced worldwide. Thus it will be helpful to the farmers in the means of cost effective farming as well as the buyers who will be consuming healthy vegetables. Thus the above mentioned data proves statistically and scientifically that Agrohomoepathy is the best method of farming with the evidence.

## Acknowledgement

I am thankful to Father Muller Homoeopathic Medical College and Hospital for funding my project and I am grateful to Rev. Fr Roshan Crasta – Administrator, Dr. ESJ Prabhu Kiran – Principal, Dr. Jacintha Monteiro- HOD, Department of Organon of Medicine and Homoeopathy

Philosophy, FMHMC. I also thank my fellowmates Sri Harene S, Gazal Mohammed, Bhuvan.M, Namith S.T and M.Chandini for their support. I also thank my parents, sister and brother in law for their encouragement and support throughout the research.

## Conflict of Interest

Not available

## Financial Support

Not available

## References

- Pereira MM, Martins AD, Morais LC, Dória J, Cavalcanti VP, Rodrigues FA, *et al.* The Potential of Agro-homeopathy Applied to Medicinal Plants-A Review. *J Agric Sci.* 2019;11(4):215-27.
- Bassi JA, Dugje IY. Effect of cowpea (*Vigna unguiculata* L. walp) in mixture with pearl millet [*Pennisetum glaucum* (L.) R. Br.] as affected by variety and time of cowpea introduction in Maiduguri North Eastern, Nigeria. *Int. J Agric. Nutr.* 2020;2(1):01-07. DOI: 10.33545/26646064.2020.v2.i1a.23
- Hahnemann S, Boericke W. *Organon of Medicine*. 6th edn. New Delhi: B. Jain; c2016.
- Hahnemann S, Hempel CJ. *Materia medica pura*. New York: William Radde.
- Cowpea. Wikipedia. Wikimedia Foundation; c2022. Available at: <https://en.wikipedia.org/wiki/Cowpea> (Accessed: December 19, 2022).
- Cowpea (no date) Cowpea | Diseases and Pests, Description, Uses, Propagation. Available at: <https://plantvillage.psu.edu/topics/cowpea/infos> (Accessed: December 19, 2022).
- Crothers L. Cowpea, Sustainable Agriculture Research & Education Program; c2021. Available at: <https://sarep.ucdavis.edu/covercrop/cowpea> (Accessed: December 19, 2022).
- Homoeopathic materia medica - by William Boericke. Available at: <http://homeoint.org/books/boericmm/> (Accessed: December 19, 2022).
- Timko MP, Ehlers JD, Roberts PA. Cowpea. In *Pulses, sugar and tuber crops*,. Springer, Berlin, Heidelberg; c2007. p. 49-67.
- Moreno NM. Agro-Homeopathy–An Alternative for Agriculture. *Hpathy Ezine*, December; c2008.
- Mandal PP, Mandal B. *Text-book of homoeopathic pharmacy*. Calcutta: New Central Book Agency; c1994.
- Mangal SK. *DNA markers in plant improvement*. New Delhi: Gene-Tech Books; c2012.
- Pharmaceutical science in homoeopathy and Pharmacodynamics. B Jain Pub Pvt Ltd; c2001.
- Soil science (no date) Welcome to Krishi Vigyan Kendra – Dakshina Kannada. Available at: [https://www.kvkdk.org/soil\\_science.html](https://www.kvkdk.org/soil_science.html) (Accessed: December 8, 2022, 14
- Systemic Agro-Homeopathy: A New Approach to Agriculture” (no date). Available at: <https://www.lidsen.com/journals/icm/icm-06-03-020>.
- RSPC. *Soils of Karnataka: Their kinds, distribution, characterization, and interpretations for optimising land use: B. executive summary*. Nagpur: National Bureau of Soil Survey and Land Use Planning, Indian Council of Agricultural Research in Barton, B. and Peat, J



- (2014). Cooperation with State Soil Survey, Dept. of Agriculture, Govt. of Karnataka; c1998.
17. Kothari CR, Garg G. Medical statistics: A guide to SPSS, Data Analysis and critical appraisal. Oxford: Wiley-Blackwell. Research methodology: Methods and techniques. New Delhi: New Age International (P) Limited, Publishers; c2019.
  18. Agro Repertory (no date). Available at: <https://homeopathyplus.com/agrorepertory.pdf>.
  19. Choudhary CM. Research methodology. Jaipur: RBSA Publishers; c2009.
  20. Author links open overlay panel Salim Laminia Person Envelope Francis Kusia Eric William Corneliusb Agyemang Danquahb Patrick Attamaha Zakaria Mukhtarua Frederick Justice Awukua Emmanuel Yaw Owusua Mavis Acheampong Gloria Mensaha *et al.* Identification of sources of resistance in cowpea lines to macrophomina root rot disease in northern Ghana, Heliyon. Elsevier; c2022. Available at: <https://www.sciencedirect.com/science/article/pii/S2405844022035058> (Accessed: December 19, 2022).
  21. Singh BB. Cowpea [*Vigna unguiculata* (L.) Walp.]. Genetic resources, chromosome engineering and crop improvement. 2005 Mar 16;1:117-62.
  22. Siddhuraju P, Becker K. The antioxidant and free radical scavenging activities of processed cowpea (*Vigna unguiculata* (L.) Walp.) seed extracts. Food chemistry. 2007 Jan 1;101(1):10-9.
  23. Hahnemann S, Boericke W. Organon of Medicine. 5th and 6<sup>th</sup>. New Delhi: B. Jain; c2016.

#### How to Cite This Article

Karanik AN, Sneha Jnanakshee HR, Kumar SS. To determine the potential of agrohomoepathy by using calcarea phosphorica 200c by seed priming technique and analyzing the germination rate and growth rate of *Vigna unguiculata* plant (cow pea) plant. International Journal of Homoeopathic Sciences. 2023;7(4):414-422.

#### Creative Commons (CC) License

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 International (CC BY-NC-SA 4.0) License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.