E-ISSN: 2616-4493
P-ISSN: 2616-4485
www.homoeopathicjournal.com
IJHS 2024; 8(4): 147-149
Received: 14-07-2024
Accepted: 23-08-2024

Dr. Kanika Kaur
Associate Professor, HOD - Department of Pathology, Shree Sardar Patel Mahila Homoeopathic Medical College & Hospital, Rajkot, Gujarat, India

Corresponding Author:
Dr. Kanika Kaur
Associate Professor, HOD - Department of Pathology, Shree Sardar Patel Mahila Homoeopathic Medical College & Hospital, Rajkot, Gujarat, India

Integrative approaches to attention deficit-hyperactivity disorder: Exploring the role of homeopathy

Dr. Kanika Kaur

DOI: https://doi.org/10.33545/26164485.2024.v8.i4c.1278

Abstract
Attention Deficit-Hyperactivity Disorder (ADHD) is a neurodevelopmental disorder that significantly affects children's capacity to function optimally across various domains of life. Individuals with ADHD commonly demonstrate developmentally inappropriate levels of inattention, hyperactivity, and impulsivity. The onset of symptoms typically occurs in early childhood and encompasses a range of challenges, including difficulties in maintaining attention, sustaining concentration, organizing tasks, completing assignments, and managing forgetfulness. Additionally, individuals may frequently misplace items, further impacting their daily functioning. The pervasive nature of these symptoms can hinder academic performance, social interactions, and overall quality of life, necessitating a comprehensive approach to diagnosis and intervention.

Keywords: ADHD, attention deficit-hyperactivity disorder, DSM-IV, DSM-V

Introduction
Attention Deficit-Hyperactivity Disorder (ADHD) is not a contemporary phenomenon; it has been recognized under various terminologies throughout history. In the 1930s, it was labeled "minimal brain dysfunction," evolving over the decades into Attention Deficit Disorder (ADD) and ultimately ADHD. Over the years, the prevalence of this disorder has increased, particularly noted during the 1950s when educational systems began to adopt more standardized practices [1].

When evaluating ADHD, it is imperative to consider what is developmentally and culturally appropriate for the individual. The disorder is conceptualized as a dysfunction of executive functioning, primarily associated with the frontal lobe. As a result, individuals with ADHD often struggle not only with attention and focus but also with decision-making and emotional regulation. Children diagnosed with ADHD frequently encounter challenges in social interactions, experience heightened frustration levels, and may engage in impulsive behaviors, often resulting in them being labeled as "troublemakers" [1].

For a diagnosis of ADHD, the symptoms must manifest before the age of 12, persist for a minimum of six months, and disrupt daily functioning. These symptoms should be observable in multiple settings, such as home, school, or during extracurricular activities. The impact of ADHD can be significant, adversely affecting social relationships, increasing risk-taking behaviors, leading to job loss, and hindering academic performance [1, 2].

Early diagnosis and intervention are crucial to mitigating the persistence of symptoms into adulthood and preventing the development of comorbid conditions. Treatment typically involves a combination of stimulant medications and psychotherapy. This review aims to offer deeper insights into the causal factors, underlying pathophysiology, and management strategies associated with ADHD [1].

Aetiology
The etiology of Attention Deficit-Hyperactivity Disorder (ADHD) is multifaceted, involving an interplay of genetic and environmental factors. ADHD is recognized as one of the most heritable psychiatric disorders, with a notably higher concordance rate observed in monozygotic twins compared to dizygotic twins, underscoring the significance of genetic predisposition. Additionally, siblings of individuals diagnosed with ADHD have a twofold increased risk of developing the disorder relative to the general population [1].

Beyond genetic influences, various environmental factors have been explored as potential contributors to the onset of ADHD. These include prenatal exposures such as viral infections, maternal smoking during pregnancy, nutritional deficiencies, and fetal alcohol exposure. Despite these investigations, brain imaging studies have not consistently yielded specific neuroanatomical findings unique to individuals with ADHD. However, existing research has revealed a reduction in dopaminergic receptors within the frontal lobes of those affected by the disorder, which may play a crucial role in its pathophysiology. Furthermore, there is emerging evidence indicating that noradrenergic receptors may also be implicated in the development of ADHD, suggesting a complex interplay of neurotransmitter systems in the disorder's manifestation [1].

Clinical Features

Inattentiveness, impulsive behavior, and hyperactivity are common developmental behaviors observed in children and adolescents. However, in the context of Attention Deficit-Hyperactivity Disorder (ADHD), these behaviors are markedly more pronounced and distinctly differ from those exhibited by their peers of the same age [3].

Children who are inattentive may struggle to listen during lessons, become easily distracted, and frequently make careless errors in their work. They often find it challenging to maintain focus during leisure activities and may exhibit forgetfulness, leading to misplaced belongings. Impulsive children may find it difficult to wait for their turn in classroom settings or during games, often interrupting others or intruding into conversations. Hyperactive children typically display fidgety behavior, such as bouncing in their seats or exhibiting difficulty remaining quiet. They may be restlessly engaged in activities like running around or climbing on furniture. The manifestations of inattentive and impulsive behavior can be exacerbated in stimulating environments that require extensive processing, such as during social gatherings or vacations [3].

The severity of ADHD and the predominance of specific behaviors can vary significantly between boys and girls, as well as among individual children. Some children may primarily exhibit inattentiveness, often appearing lost in thought or daydreaming, while others may demonstrate more pronounced impulsivity and hyperactivity. Based on which behaviors are most evident, children with ADHD are typically classified into two subtypes: primarily inattentive or primarily hyperactive-impulsive. Those exhibiting significant inattentiveness without hyperactivity are often referred to as having Attention Deficit Disorder (ADD) [3].

Diagnosis

Attention Deficit-Hyperactivity Disorder (ADHD) is primarily diagnosed through a comprehensive clinical evaluation rather than relying on specific laboratory or radiological tests. Neuropsychological assessments, while informative, do not possess sufficient sensitivity to serve as standalone diagnostic tools for ADHD; thus, the diagnosis is predominantly based on a detailed patient history [1].

The evaluation process typically involves utilizing various standardized rating scales and collecting information from multiple informants, including parents and teachers, to gain a holistic view of the child's behavior across different environments. Additionally, clinicians must remain vigilant for the presence of comorbid disorders that may elucidate or overlap with the symptoms exhibited by the child. ADHD should not be diagnosed if the observed symptoms can be more accurately attributed to another condition, such as a psychotic disorder or a manic episode, thereby ensuring a precise and accurate clinical assessment [1].

DSM 5: Types of ADHD

Predominantly inattentive.
Predominantly impulsive or hyperactive.
Combination of the above.
The onset is usually before age 12.
Symptoms present at school, work, or home.
The disturbance causes significant impairment in social, occupational, and academic functioning.
The disorder is not accounted for by any other behavior disorder.

Differential Diagnosis

Differentiating Attention Deficit-Hyperactivity Disorder (ADHD) from other clinical disorders is essential, given the significant overlap in symptoms among various conditions. Mood disorders, including depression and anxiety, can often be misdiagnosed in individuals with ADHD, as both conditions may present with symptoms such as inattention, poor concentration, memory deficits, and distractibility [1].

Moreover, the relationship between ADHD and substance abuse disorders requires careful consideration, as children diagnosed with ADHD are at an elevated risk for developing substance use issues later in life [1].

When diagnosing ADHD, it is imperative to thoroughly evaluate and rule out other potential contributing factors, including hearing disorders, learning disabilities, and developmental disorders. This comprehensive approach ensures an accurate assessment of the child's condition, facilitating the implementation of appropriate interventions and treatment strategies tailored to their specific needs. By addressing these comorbidities and differential diagnoses, clinicians can provide a more effective and holistic care plan for individuals exhibiting symptoms of ADHD [1].

Treatment/management

Pharmacological therapy serves as the cornerstone of treatment for individuals diagnosed with Attention Deficit-Hyperactivity Disorder (ADHD), encompassing two principal categories: stimulants and non-stimulants. Stimulant medications are the most widely utilized and are subdivided into two primary classifications: amphetamines and methylphenidates. These medications are known for their efficacy in enhancing attention and reducing hyperactive and impulsive behaviors [1, 4].

On the other hand, non-stimulant options for managing ADHD include two key types: antidepressants and alpha-agonists. Antidepressants, such as certain selective serotonin reuptake inhibitors (SSRIs) and tricyclic antidepressants, can be beneficial in addressing ADHD symptoms, particularly in cases where stimulants are contraindicated or poorly tolerated. Alpha agonists, such as guanfacine and clonidine, are also employed to manage symptoms, particularly in situations where behavioral issues are prominent or when there is a need to address comorbid conditions like anxiety or sleep disturbances. Overall, the selection of pharmacological therapy should be tailored to the individual patient's needs, considering factors such as symptom profile, comorbidities, and potential side effects, to achieve optimal therapeutic outcomes [1, 4].

Homoeopathic management

1. **Belladonna:** Lives in a world of their own, focused on visions and specters, detached from reality. The retina is unresponsive to real objects, but hallucinations and vivid illusions dominate their mind. Engaged in a chaotic world of visual hallucinations, often seeing monsters and frightening faces. Experiences delirium with violent, aggressive behavior and a desire to escape. May lose consciousness, become emotionally unstable, and show heightened sensitivity to all senses [5].
2. **Causticum:** A child who fears going to bed alone, easily brought to tears by even minor things. Experiences deep sadness and hopelessness and is highly sympathetic. Symptoms worsen when thinking about their ailments, particularly with long-term grief or sudden emotional upheavals [5].
3. **Hyoscyamus:** Very distrustful and suspicious. Becomes overly talkative, often displaying inappropriate or obscene behavior. Manic, revealing their body or acting jealously and foolishly. Excessive laughter, even at inappropriate times. Delirium can lead to attempts to escape, with low, incoherent speech and deep stupor [5].
4. **Lycopodium:** Prone to sadness, especially fear of being alone. Easily irritated by minor issues and highly sensitive. Hesitant to start new tasks, stubborn and arrogant when ill. Lacks self-confidence and rushes through meals. Feels overwhelmed by stress and apprehensive, with poor memory and mental confusion. May write or spell incorrectly and have a declining ability to think clearly. Cannot tolerate new things or read their own writing. Often wakes up feeling sad in the morning [5].
5. **Natrum muriaticum:** Symptoms stem from emotional causes like grief, fear, or anger. Prone to depression, especially in chronic illnesses, and worsened by consolation. Easily irritated by small matters and tends to act awkward or hasty. Prefers solitude to cry and experiences tears even while laughing [5].
6. **Nux vomica:** Highly irritable and sensitive to all stimuli. Becomes unpleasant, malicious, and easily disturbed by noises, smells, and light. Dislikes being touched and feels that time moves too slowly. Small ailments have a significant impact, leading to resentment and blame towards others. Tends to be sullen and overly critical [5].
7. **Pulsatilla:** Tends to cry easily and is timid and indecisive. Fears being alone in the evening, the dark, or seeing ghosts. Seeks sympathy and affection, especially from children. Easily discouraged, with a heightened fear of the opposite sex and religious melancholy. Mood swings frequently, alternating between joy and sorrow [5].
8. **Phosphorus:** Experiences deep depression and irritability. Fearful, as if something is lurking everywhere. Tends to be clairvoyant and easily startled. Extremely sensitive to external stimuli, with memory loss and potential insanity. Feels euphoric and has a heightened sense of self-importance. Restless and fidgety, indifferent to emotions. Afraid of being alone, with a constant dread of death. The brain feels fatigued [5].
9. **Sulphur:** Extremely forgetful and has difficulty thinking clearly. Prone to delusions, believing that ragged things are beautiful or that they are immensely wealthy. Always busy but irritable and selfish, with no regard for others. Experiences religious melancholy and a lack of motivation for work. Imagines giving harmful things to people, leading to their death. Typically irritable, depressed, thin, and weak, even with a good appetite [5].

References

1. Magnus W, Nazir S, Anilkumar AC, Shaban K. Attention Deficit Hyperactivity Disorder. In: StatPearls. Treasure Island (FL): StatPearls Publishing; 2024.
2. Maitre S. Attention Deficit Hyperactivity Disorder in Childhood: Overview, Diagnosis and Treatment. Am Med Assoc J Ethic. 2007;9(6):433-436.
3. Institute for Quality and Efficiency in Health Care (IQWiG). Overview: Attention deficit hyperactivity disorder (ADHD); 2022.
4. Drechsler R, Brem S, Brandeis D, Grünblatt E, Berger G, Walitza S *et al*. ADHD: Current concepts and treatments in children and adolescents. Neuropediatrics. 2020;51(5):315-335.
5. Boericke W. Pocket manual of homoeopathic materia medica. 9th ed. New Delhi: IBPP; 2015. p. 100, 161, 297, 362, 408, 422, 450, 478, 545.

How to Cite This Article
Kaur K. Integrative approaches to attention deficit-hyperactivity disorder: Exploring the role of homeopathy. International Journal of Homoeopathic Sciences 2024; 8(4): 147-149.