International Journal of Homoeopathic Sciences

E-ISSN: 2616-4493
P-ISSN: 2616-4485
www.homoeopathicjournal.com
IJHS 2025; 9(1): 757-760
Received: 23-12-2024
Accepted: 29-01-2025

Dr. Avani Roy
MD Scholar, Department of Psychiatry, Government Homoeopathic Medical College & Hospital, Bhopal, Madhya Pradesh, India

Dr. Santosh Hande
HOD, Department of Psychiatry, Government Homoeopathic Medical College & Hospital, Bhopal, Madhya Pradesh, India

Corresponding Author:
Dr. Avani Roy
MD Scholar, Department of Psychiatry, Government Homoeopathic Medical College & Hospital, Bhopal, Madhya Pradesh, India

Holistic approaches in the treatment of generalized anxiety disorder in adolescents: A focus on homoeopathy and therapy

Avani Roy and Santosh Hande

DOI: https://www.doi.org/10.33545/26164485.2025.v9.i1.K.1433

Abstract
Generalized Anxiety Disorder (GAD) is a prevalent and concerning mental health condition affecting adolescents globally. With its insidious onset and wide-ranging impact, GAD can lead to severe long-term effects, including academic struggles, social withdrawal, substance abuse, and in some cases, suicidal tendencies. This article explores the various causes and risk factors of GAD in adolescents, its epidemiology, and its significant impact on adolescent development. It further delves into the biopsychosocial model of the disorder and the most effective management strategies, including both conventional therapeutic options and the emerging role of alternative treatments, particularly homeopathy. As the prevalence of anxiety disorders continues to rise among adolescents, understanding the multifactorial nature of this condition is essential for developing holistic, individualized care and intervention strategies.

Keywords: Generalized Anxiety Disorder (GAD), adolescents, holistic approaches

Introduction
Generalized Anxiety Disorder (GAD) in adolescents involves feeling excessively worried or tense about everyday situations, even when there's no real reason to be anxious. This worry lasts for at least six months and makes it hard for the young person to carry out daily activities [1]. Adolescence is a time when young people go through many changes, both physically and emotionally, and it typically refers to anyone between the ages of 10 and 19, according to the World Health Organization (WHO). During this time, adolescents are learning how to handle their emotions and cope with life's challenges. For those with anxiety, a lot of their worry is focused on things like school performance and whether they are good enough in different areas of their lives [2].
The prevalence of GAD among adolescents is on the rise, with various genetic, neurobiological, psychological, and environmental factors contributing to its onset. The World Health Organization (WHO) classifies anxiety disorders as one of the leading causes of illness among adolescents, with GAD being the fourth leading cause of disability in this age group. As the understanding of GAD deepens, more emphasis is being placed on early detection, individualized treatment plans, and the integration of holistic therapeutic approaches [3].

Epidemiology
Generalized Anxiety Disorder is a common psychiatric condition among adolescents, with studies showing that approximately 31.9% of adolescents experience some form of anxiety disorder, with 8.3% of these individuals experiencing severe mental impairment due to anxiety. The prevalence of GAD is higher among older adolescents, with around 4.6% of adolescents aged 15 to 19 and 3.6% of those aged 10 to 14 experiencing the disorder. In terms of gender, anxiety disorders tend to affect females more than males, with a ratio of 2:1 in adolescence [4].

Etiology
GAD in adolescents arises from the interplay of various biological, psychological, and environmental factors. These factors contribute to the heightened vulnerability of adolescents

to anxiety disorders, making this a multifaceted condition that requires a comprehensive understanding for effective management.

1. **Biological Factors**

- **Genetic factor:** The development of GAD has a strong genetic component, with studies indicating that individuals with a family history of anxiety or mood disorders are more likely to develop GAD themselves. Genetic predisposition plays a role in the increased susceptibility of adolescents to anxiety disorders, with twin studies suggesting that the heritability of anxiety disorders may range from 30% to 50% [5].
- **Neurobiological mechanism:** Neurobiologically, dysfunctions in brain areas such as the amygdala, prefrontal cortex, and hippocampus contribute to heightened anxiety. These areas are critical for processing emotional stimuli, regulating fear responses, and controlling stress. Adolescents with GAD often exhibit an overactive hypothalamic-pituitary-adrenal (HPA) axis, leading to an exaggerated stress response [6]. Furthermore, imbalances in neurotransmitters such as serotonin, gamma-aminobutyric acid (GABA), and norepinephrine are also implicated in the development of GAD. These imbalances disrupt the brain's ability to regulate emotions, leading to chronic anxiety symptoms [7].

2. **Psychological Factors**

- **Cognitive Distortion:** Adolescents with GAD tend to have specific cognitive and personality traits that make them more vulnerable to developing anxiety. Cognitive distortions, such as catastrophizing, overestimating the likelihood of negative events, and underestimating one's ability to cope with stress, are common among adolescents with GAD. These maladaptive thought patterns contribute to excessive worry and a sense of impending doom [8].
- **Personality traits:** Adolescents with GAD often exhibit perfectionistic tendencies, where they set unrealistically high standards for themselves and experience intense fear of failure. This perfectionism, combined with low self-esteem, creates a cycle of anxiety that becomes difficult to break. Emotional dysregulation, a common feature in adolescence due to hormonal changes and identity formation, further exacerbates anxiety symptoms [9].

3. **Environmental Factors**

- **Family dynamics:** Several external stressors significantly contribute to the development of GAD in adolescents. Family dynamics, such as overprotective or critical parenting styles, contribute to emotional instability and a lack of coping mechanisms. Adolescents who experience familial stressors like parental separation or trauma are at an increased risk of developing GAD [10].
- **Environmental factors:** Social stressors are also key contributors. Academic pressures, bullying, peer rejection, and social comparison, often amplified by social media, create a perfect storm for anxiety in adolescents. The societal pressure to succeed academically and socially can be overwhelming, and the constant need for validation, especially in the age of social media, significantly contributes to anxiety and worry about self-image and social acceptance [11].

Impact of Generalized Anxiety Disorder on Adolescents

The effects of GAD on adolescents are far-reaching and can have long-lasting consequences on their development. The impact is most noticeable in academic, social, and emotional spheres:

1. **Academic Impairment**

Anxiety often manifests in poor concentration, fatigue, and a constant sense of dread about school performance. Adolescents with GAD may become overwhelmed by the pressure to succeed, leading to a decline in academic performance. This can create a cycle of failure and worry, further exacerbating the anxiety [12].

2. **Social Withdrawal**

Adolescents with GAD often retreat from social interactions due to fears of rejection or not meeting social expectations. They may avoid activities such as group work or extracurricular activities, which limits their opportunities for social engagement and emotional growth [13].

3. **Comorbid Conditions**

GAD often co-occurs with other mental health disorders such as depression, substance abuse, and eating disorders. The constant anxiety and worry may lead to emotional exhaustion and feelings of helplessness, triggering depressive symptoms. Additionally, adolescents may turn to substances as a coping mechanism, further complicating their mental health [14].

4. **Increased Risk of Suicide**

The most alarming consequence of untreated GAD is the increased risk of suicide. Adolescents with chronic anxiety are at a higher risk of developing suicidal thoughts and behaviors. According to the WHO, suicide is the third leading cause of death among individuals aged 15-29, with anxiety disorders being a significant contributing factor [15].

Diagnosis

DSM-V Diagnostic Criteria for Generalized Anxiety Disorder (GAD)

Clinicians assess GAD by examining the following criteria:

A. Excessive anxiety and worry about various subjects, events, or activities. Worry occurs more often for at least 6 months.
B. Worry is considered very difficult to manage. The concerns of both adults and children can easily shift from one topic to another.
C. At least three of the following physical or cognitive symptoms are associated with anxiety and worry (only one symptom is needed to diagnose GAD in children):

- Agitation or restlessness
- Tires easily; more tired than usual
- Decreased ability to concentrate or feeling like your mind is going blank
- Irritability (which may or may not be visible to others)
- Increased muscle pain or tenderness
- Sleep disturbances (difficulty falling or staying asleep, restlessness at night, or unsatisfactory sleep)

D. The anxiety, worry, or physical symptoms cause clinically significant distress in social, occupational, or other areas of functioning.
E. The disturbance is not related to the physiological effects of a substance or any other medical condition.
F. The disturbance is not better explained by another medical condition [16].

ICD-10 Classification for Generalized Anxiety Disorder Code: F41.1

Diagnostic Criteria

The patients with Generalized Anxiety Disorder (GAD) typically experience primary symptoms of anxiety most days for a duration of several weeks, and often extending over several months. These symptoms are manifested through the following:

- **Apprehension:** Persistent worries about the future, fears of misfortune, feeling "on edge," difficulty concentrating, and an overall sense of unease.
- **Motor Tension:** Physical manifestations such as restlessness, fidgeting, tension headaches, trembling, and an inability to relax.
- **Autonomic Overactivity:** Symptoms like lightheadedness, sweating, rapid heart rate (tachycardia), shortness of breath (tachypnea), epigastric discomfort, dizziness, dry mouth, and other physical signs of stress [17].

Management of Generalized Anxiety Disorder in Adolescents

The treatment of GAD in adolescents typically involves a combination of therapeutic approaches aimed at addressing the disorder's underlying causes and mitigating its symptoms.

1. Cognitive Behavioral Therapy (CBT)

Cognitive Behavioral Therapy is considered the gold standard in treating GAD in adolescents. CBT helps adolescents identify and challenge maladaptive thought patterns and beliefs, replacing them with more adaptive coping mechanisms. It also encourages gradual exposure to anxiety-provoking situations, reducing avoidance behaviors. Studies have shown that CBT can significantly reduce anxiety symptoms and improve functioning in adolescents [18].

2. Mindfulness-Based Stress Reduction (MBSR)

Mindfulness-based therapies help adolescents cultivate awareness of the present moment, which can counteract the tendency to worry about the future. Techniques such as deep breathing, body scanning, and meditation can help break the cycle of anxiety and promote emotional regulation. MBSR has been shown to reduce anxiety and improve well-being in adolescents with GAD [19].

3. Family Therapy

Family therapy can be beneficial in addressing the familial dynamics that contribute to the adolescent's anxiety. This approach helps family members improve communication, resolve conflicts, and support the adolescent's emotional needs [20].

Homeopathic Treatment

Homeopathy offers a holistic approach to managing GAD, focusing on individualized remedies based on the adolescent's emotional and physical symptoms. The goal of homeopathy is to restore balance to the body's natural processes, addressing both the root causes of anxiety and the manifestation of symptoms.

Aconitum napellus

This remedy is indicated for adolescents with sudden, intense anxiety often following a shock or traumatic event. These individuals may experience restlessness, fear of impending doom, and heightened panic. Symptoms include agitation, palpitations, shortness of breath, or a choking sensation. Anxiety can worsen in the evening or night, causing insomnia and nightmares. Adolescents may fear being alone and seek reassurance. They are hypersensitive to noise, light, or touch and feel overwhelmed by their surroundings, sometimes exhibiting a sense of disconnection from reality.

Argentum nitricum

Adolescents who need this remedy often experience anticipatory anxiety, particularly regarding upcoming events. They are restless, irritable, and impulsive, exhibiting inner turmoil. Physical symptoms include trembling, palpitations, and gastrointestinal issues like diarrhoea. Anxiety is often rooted in a fear of failure, performance anxiety, and a constant need for reassurance. There is a compulsive urge to act quickly, leading to hasty decisions and impulsive behaviours.

Arsenicum album

Adolescents needing this remedy may experience profound anxiety marked by insecurity and a fear of death or illness. These individuals exhibit excessive worry about health and safety, feel exhausted yet cannot relax, and may engage in compulsive behaviors to alleviate their fears. Physical symptoms include insomnia, digestive issues, and cold extremities. They tend to seek company but fear being left alone, exhibiting a paradoxical desire for companionship. Anxiety intensifies at night, disrupting sleep.

Calcarea carbonica

Adolescents who benefit from this remedy typically experience anxiety tied to a sense of inadequacy and being overwhelmed by daily responsibilities. There is a fear of failure, often accompanied by physical symptoms like palpitations, sweating, and digestive disturbances. Anxiety is linked to a need for stability, making them fearful of changes or uncertainties. These individuals tend to overthink situations and may be indecisive. They may have specific cravings, such as for eggs, milk, or sweets, and may fear heights or enclosed spaces.

Lycopodium

Adolescents requiring this remedy typically experience anxiety linked to a fear of failure and self-doubt. They may feel insecure and perfectionistic, leading to significant anxiety about performance and decision-making. Physical symptoms often include digestive disturbances and palpitations. There is a fear of social situations, and they tend to feel more comfortable in familiar environments. Anxiety is exacerbated by responsibilities and a fear of being judged or criticized, both in personal and professional contexts.

Phosphorus

Adolescents needing this remedy often experience anxiety linked to sensitivity and emotional overwhelm. They may fear abandonment and have a deep need for affection and reassurance. Anxiety can manifest through restlessness, insomnia, and physical symptoms like palpitations and digestive disturbances. These individuals are highly influenced by others' emotions and may feel easily drained by social interactions. They may also experience vivid

dreams and nightmares, exacerbating their anxiety.

Pulsatilla
This remedy is suitable for adolescents with anxiety marked by emotional instability and sensitivity. They may experience a fear of abandonment and constantly seek reassurance from others. Hormonal changes or emotional turmoil often trigger their anxiety, resulting in physical symptoms like headaches, digestive issues, and a sense of heaviness. They feel better in the open air, may crave company, and experience feelings of loneliness or neglect. These individuals are often indecisive and changeable, reflecting their fluctuating emotions [21].

Conclusion
Generalized Anxiety Disorder (GAD) in adolescents is a critical mental health concern that requires effective, holistic treatment strategies. While conventional therapies such as Cognitive Behavioural Therapy (CBT) and pharmacotherapy are widely recognized, homeopathy offers a valuable, individualized approach. By addressing the emotional, psychological, and physical symptoms specific to each adolescent, homeopathic remedies such as Aconitum Napellus, Argentum Nitricum, and Pulsatilla etc. can provide safe, non-invasive relief. The holistic nature of homeopathy supports the body's natural healing processes and restores balance, making it an effective complementary treatment for GAD. As anxiety disorders continue to rise, integrating homeopathy into a comprehensive treatment plan can offer significant benefits for adolescents, enhancing their overall well-being and mental health.

Conflict of Interest
Not available

Financial Support
Not available

References
1. Gale CK, Millichamp J. Generalised anxiety disorder in children and adolescents. BMJ Clin Evid. 2016 Jan 13;2016:1002. PMID: 26763675; PMCID: PMC4711893.
2. Adolescence. Britannica. Available from: http://www.britanica.com/science/adolescence.
3. Adolescent health. World Health Organization. Available from: http://www.who.int/news-room/fact-sheets/detail/adolescent.
4. Garcia I, O'Neil J. Anxiety in adolescents. J Nurse Pract. 2021;17(1):49-53. ISSN 1555-4155.
5. Hettema JM, Neale MC, Kendler KS. A review and meta-analysis of the genetic epidemiology of anxiety disorders. Am J Psychiatry. 2001;158(10):1568-78.
6. Beesdo K, Knappe S, Pine DS. Anxiety and anxiety disorders in children and adolescents: Developmental issues and implications for DSM-V. Psychiatr Clin North Am. 2009;32(3):483-524.
7. Mathew SJ, Price RB, Charney DS. Recent advances in the neurobiology of anxiety disorders: Implications for novel therapeutics. Am J Med Genet C Semin Med Genet. 2008;148C(2):89-98.
8. Hirsch CR, Mathews A. A cognitive model of pathological worry. Behav Res Ther. 2012;50(10):636-46.
9. Zinbarg RE, Mineka S, Bobova L, Craske MG, Vrshek-Schallhorn S, Griffith JW, *et al*. Testing a hierarchical model of neuroticism and its cognitive facets: Prospective relations to anxiety and depression in two samples. J Abnorm Psychol. 2016;125(6):798-819.
10. Ginsburg GS, Schlossberg MC. Family-based treatment of childhood anxiety disorders. Int Rev Psychiatry. 2002;14(2):143-54.
11. Twenge JM, Campbell WK. Media use is linked to lower psychological well-being: Evidence from three datasets. Psychol Pop Media Cult. 2019;8(4):491-501.
12. Masi G, Perugi G, Giannotti C, *et al*. Anxiety disorders and academic performance in children and adolescents: A systematic review. Child Psychiatry Hum Dev. 2014;45(5):575-583.
13. Beidel DC, Turner SM. Social phobia and social withdrawal in children and adolescents. J Clin Child Psychol. 2007;36(3):410-418.
14. Kessler RC, Chiu WT, Demler O, *et al*. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. Arch Gen Psychiatry. 2005;62(6):617-627.
15. World Health Organization (WHO). Suicide prevention. Geneva: WHO; 2014. Available from: https://www.who.int/mental_health/prevention/suicide/suicideprevent/en/
16. American Psychiatric Association. Diagnostic and statistical manual of mental disorders. BMC Med. 2013;17:133-7.
17. World Health Organization. International Classification of Diseases 11th Revision (ICD-11). Generalized Anxiety Disorder. Geneva: WHO; 2019. Available from: https://icd.who.int/browse11/l-m/en
18. Piacentini J, Peris TS, Bergman RL, Chang S, Jaffer M. Functional impairment in childhood OCD: Development and psychometric properties of the Child Obsessive-Compulsive Impact Scale-Revised (COIS-R). J Clin Child Adolesc Psychol. 2007;36(4):645-653. doi:10.1080/15374410701662790.
19. Biegel GM, Brown KW, Shapiro SL, Schubert CM. Mindfulness-based stress reduction for the treatment of adolescent psychiatric outpatients: A randomized clinical trial. J Consult Clin Psychol. 2009;77(5):855-866. doi:10.1037/a0016241.
20. Wood JJ, McLeod BD, Piacentini JC, Sigman M. One-year follow-up of family versus child CBT for anxiety disorders: Exploring the roles of child age and parental intrusiveness. Child Psychiatry Hum Dev. 2009;40(3):301-316. doi:10.1007/s10578-009-0123-5.
21. Kent JT. Lectures on Homoeopathic Materia Medica. 2nd ed. New Delhi: B. Jain Publishers; c2003.

How to Cite This Article
Roy A, Hande S. Holistic approaches in the treatment of generalized anxiety disorder in adolescents: A focus on homoeopathy and therapy. International Journal of Homoeopathic Sciences. 2025;9(1):757-760.